

Foundations of Privacy 2015

Recitation on Logic

Amit Datta



Blackboard Issues?

Propositional (“0th order”) Logic

Propositional (“0th order”) Logic

A model for a simple **subset**
of mathematical reasoning

Not	\neg
And	\wedge
Or	\vee
Implies	\rightarrow
If And Only If	\leftrightarrow

Propositional (“0th order”) Logic

An English statement that can be true or false

Propositional variable:
a symbol (letter) representing it

“Potassium is observed.”

k

“Hydrogen is observed.”

h

“Pixel 29 is black.”

p_{29}

“It’s raining.”

r

Compound sentence

Potassium is not observed.

At least one of hydrogen and potassium is observed.

If potassium is observed then hydrogen is also observed.

If I'm not playing tennis then I'm watching tennis, and if I'm not watching tennis then I'm reading about tennis.

Propositional formula

$\neg k$

$(h \vee k)$

$(k \rightarrow h)$

p, w, r

$((\neg p \rightarrow w) \wedge (\neg w \rightarrow r))$

Formally, formulas are strings made up of:

((punctuation)
)	(punctuation)
\neg	(not)
\wedge	(and)
\vee	(or)
\rightarrow	(implies)
\leftrightarrow	(if and only if)
x_1, x_2, x_3, \dots	(variable symbols)

Well-formed formula (WFF)

= A string which is syntactically “legitimate”.

WFF

x_1

$((x_1 \wedge (x_3 \rightarrow \neg x_2)) \vee x_1)$

$\neg((x_{10} \leftrightarrow x_{11}) \wedge (x_2 \rightarrow x_5))$

not a WFF

$x_1 \wedge$

$)x_2 \rightarrow \rightarrow$

$((x_1 \wedge (x_3 \rightarrow \neg x_2)) \neg x_1)$

Well-formed formula (WFF)

Formally, WFFs have an inductive definition:

Base case:

Single variables are WFFs.

Inductive rules:

- If A is a WFF, so is $\neg A$.
- If A, B are WFFs, so are
 - $(A \wedge B)$,
 - $(A \vee B)$,
 - $(A \rightarrow B)$,
 - $(A \leftrightarrow B)$.



Let's talk about **TRUTH (SEMANTICS)**.

“If potassium is observed then carbon and hydrogen are also observed.”

$(k \rightarrow (c \wedge h))$

Q: Is this statement true?

A: The question does not make sense.

“If potassium is observed then carbon and hydrogen are also observed.”

$$(k \rightarrow (c \wedge h))$$

Whether this statement/formula is true/false depends on whether the variables are true/false (“state of the world”).

If k is T, c is T, h is F...
... the formula is **False**.

If k is F, c is F, h is T...
... the formula is **True**.

Truth assignment V :

assigns T or F to each variable

Extends to give a **truth value $V[S]$**
for any formula S by applying these rules:

A	B	$\neg A$	$(A \wedge B)$	$(A \vee B)$	$(A \rightarrow B)$	$(A \leftrightarrow B)$
F	F	T	F	F	T	T
F	T	T	F	T	T	F
T	F	F	F	T	F	F
T	T	F	T	T	T	T

Truth assignment V :

assigns T or F to each variable

Extends to give a **truth value $V[S]$**
for any formula S by applying these rules:

A	B	$\neg A$	$(A \wedge B)$	$(A \vee B)$	$(A \rightarrow B)$	$(A \leftrightarrow B)$	$\neg A \vee B$
F	F	T	F	F	T	T	T
F	T	T	F	T	T	F	T
T	F	F	F	T	F	F	F
T	T	F	T	T	T	T	T

Truth assignment example

$$S = (x_1 \rightarrow (x_2 \wedge x_3))$$

$$\mathbf{V} : \begin{array}{l} x_1 = \mathbf{T} \\ x_2 = \mathbf{T} \\ x_3 = \mathbf{F} \end{array}$$

$$\mathbf{V}[S] = (\mathbf{T} \rightarrow (\mathbf{T} \wedge \mathbf{F}))$$

$$\mathbf{V}[S] = (\mathbf{T} \rightarrow \mathbf{F})$$

$$\mathbf{V}[S] = \mathbf{F}$$

Satisfiability

V satisfies S:

$$V[S] = T$$

S is satisfiable:

there exists **V** such that $V[S] = T$

S is unsatisfiable:

$$V[S] = F \text{ for all } V$$

S is a tautology:

$$V[S] = T \text{ for all } V$$

All well-formed formulas

unsatisfiable

$(k \wedge \neg k)$

satisfiable

$(k \rightarrow (c \wedge h))$

tautology

$(h \rightarrow h)$

“Potassium is observed and potassium is not observed.”

“If potassium is observed then
carbon and hydrogen are observed.”

“If hydrogen is observed then hydrogen is observed.”

Tautology: automatically true,
for 'purely logical' reasons

Unsatisfiable: automatically false,
for purely logical reasons

Satisfiable (but not a tautology):

truth value depends
on the state of the world

$$S = ((x \rightarrow (y \rightarrow z)) \leftrightarrow ((x \wedge y) \rightarrow z))$$

Truth table

x	y	z	$((x \rightarrow (y \rightarrow z)) \leftrightarrow ((x \wedge y) \rightarrow z))$
F	F	F	
F	F	T	
F	T	F	
F	T	T	
T	F	F	
T	F	T	
T	T	F	
T	T	T	

$$S = ((x \rightarrow (y \rightarrow z)) \leftrightarrow ((x \wedge y) \rightarrow z))$$

Truth table

x	y	z	$((x \rightarrow (y \rightarrow z)) \leftrightarrow ((x \wedge y) \rightarrow z))$
F	F	F	T
F	F	T	
F	T	F	
F	T	T	
T	F	F	
T	F	T	
T	T	F	
T	T	T	

S is satisfiable!

$$S = ((x \rightarrow (y \rightarrow z)) \leftrightarrow ((x \wedge y) \rightarrow z))$$

Truth table

x	y	z	$((x \rightarrow (y \rightarrow z)) \leftrightarrow ((x \wedge y) \rightarrow z))$
F	F	F	T
F	F	T	T
F	T	F	T
F	T	T	T
T	F	F	T
T	F	T	T
T	T	F	T
T	T	T	T


S is a tautology!

Problem: Show $((x \rightarrow y) \wedge x) \rightarrow y$ is a tautology.

Truth-table method: quite long, not intuitive

Can we build up a proof systematically?

Inference rules:



$$\begin{array}{c}
 \frac{(x \rightarrow y) \wedge x}{(x \rightarrow y)} \quad \wedge \text{ elimination} \qquad \frac{(x \rightarrow y) \wedge x}{x} \quad \wedge \text{ elimination} \\
 \frac{(x \rightarrow y) \quad x}{y} \quad \rightarrow \text{ elimination} \\
 \frac{y}{((x \rightarrow y) \wedge x) \rightarrow y} \quad \rightarrow \text{ introduction}
 \end{array}$$

What is a proof?

A sequence of statements,
 each of which
 is an axiom,
 or a hypothesis,
 or follows from previous statements
 using an inference rule

Problem: Show $((x \rightarrow y) \wedge x) \rightarrow y$ is a tautology.

Solution 1: Truth-table method (semantic proof)

Solution 2: Use proof system: (syntactic proof)

Are all theorems (whatever can be proved) tautology?

Yes...for propositional logic

This property is called soundness of propositional logic

Are all tautology theorems?

Yes...for propositional logic

This property is called completeness of propositional logic



First order logic (FOL)

A model for pretty much all mathematical reasoning

Not, And, Or, Implies, If And Only If

Plus: For All (\forall), There Exists (\exists), Equals (=)
“constants”, “predicates”, “functions”

Variables like x now represent **objects**, not truth-values.

“Alex is smarter than everyone”:

$\forall x \text{ IsSmarter}(a, x)$



variable:
stands for an
object (person)

constant name:
stands for a
particular object

predicate name:
stands for a mapping,
object(s) \mapsto T/F

quantifier

“Alex is smarter than everyone”:

$\forall x \text{ IsSmarter}(a, x)$

“Alex is smarter than everyone else”:

$\forall x (\neg(x=a) \rightarrow \text{IsSmarter}(a, x))$

propositional logic, as usual

equality (of objects)

“Alex is smarter than everyone”:

$\forall x \text{ IsSmarter}(a, x)$

“Alex is smarter than everyone else”:

$\forall x (\neg(x=a) \rightarrow \text{IsSmarter}(a, x))$

“Alex’s father is smarter than everyone else’s father”:

$\forall x (\neg(x=a) \rightarrow \text{IsSmarter}(\text{Father}(a), \text{Father}(x)))$

function name:

stands for a mapping,
object(s) \mapsto object

Vocabulary: A collection of constant-names,
function-names,
predicate-names.

Vocabulary from the previous slide:

one constant-name: a

one function-name: Father(·)

one predicate-name: IsSmarter(·, ·)

Vocabulary: A collection of constant-names,
function-names,
predicate-names.

Another example of a vocabulary:

one constant-name: **a**

two function-names: **Next(·), Combine(·, ·)**

one predicate-name: **IsPrior(·, ·)**

Example (well-formed) “sentences”:

$$\exists x (\text{Next}(x)=a)$$

$$\forall x \forall y (\text{IsPrior}(x, \text{Combine}(a, y)) \rightarrow (\text{Next}(x)=y))$$

$$(\forall x \text{ IsPrior}(x, \text{Next}(x))) \rightarrow (\text{Next}(a)=\text{Next}(a))$$



Let's talk about **TRUTH.**

$\exists x (\text{Next}(x) = \text{Combine}(a, a))$

Q: Is this sentence true?

A: The question does not make sense.

Whether or not this sentence is true depends on the **interpretation** of the vocabulary.

Interpretation:

Informally, says what objects are and what the vocabulary means.

$\exists x (\text{Next}(x)=\text{Combine}(a,a))$

Q: Is this sentence true?

A: The question does not make sense.

Whether or not this sentence is true depends on the **interpretation** of the vocabulary.

Interpretation:

Specifies a nonempty set (“**universe**”) of objects.

Maps each **constant-name** to a **specific object**.

Maps each **predicate-name** to an **actual predicate**.

Maps each **function-name** to an **actual function**.

$\exists x (\text{Next}(x) = \text{Combine}(a, a))$

Interpretation #1:

- Universe = all strings of 0's and 1's
- $a = 1001$
- $\text{Next}(x) = x0$
- $\text{Combine}(x, y) = xy$
- $\text{IsPrior}(x, y) = \text{True}$ iff x is a prefix of y

For this interpretation,
the sentence is...

...False

$\exists x (\text{Next}(x) = \text{Combine}(a, a))$

Interpretation #2:

- Universe = integers
- $a = 0$
- $\text{Next}(x) = x + 1$
- $\text{Combine}(x, y) = x + y$
- $\text{IsPrior}(x, y) = \text{True}$ iff $x < y$

For this interpretation,
the sentence is...

...True

$(x = 1)$

$\exists x (\text{Next}(x) = \text{Combine}(a, a))$

Interpretation #2:

- Universe = positive integers
- $a = 0$
- $\text{Next}(x) = x + 1$
- $\text{Combine}(x, y) = x + y$
- $\text{IsPrior}(x, y) = \text{True}$ iff $x < y$

For this interpretation,
the sentence is...

...**False**

Satisfiability / Tautology

Interpretation I satisfies sentence S :

$$I[S] = T$$

S is satisfiable:

there exists I such that $I[S] = T$

S is unsatisfiable:

$$I[S] = F \text{ for all } I$$

S is a tautology:

$$I[S] = T \text{ for all } I$$

All sentences in a given vocabulary

unsatisfiable

$$\exists x \neg(\text{Next}(x)=\text{Next}(x))$$

satisfiable

$$\exists x (\text{Next}(x)=\text{Combine}(a,a))$$

tautology

$$(\forall x(x=a)) \rightarrow (\text{Next}(a)=a)$$

Tautology: automatically true,
for 'purely logical' reasons

Unsatisfiable: automatically false,
for purely logical reasons

Satisfiable (but not a tautology):

truth value depends
on the interpretation
of the vocabulary

$$(\exists y \forall x (x = \text{Next}(y))) \rightarrow (\forall w \forall z (w = z))$$

Problem 1: Show this is satisfiable.

Let's pick this interpretation:

Universe = integers, $\text{Next}(y) = y+1$.

Now $(\exists y \forall x (x = \text{Next}(y)))$ means

“there's an integer y such
that every integer $= y+1$ ”.

That's False!

So the whole sentence becomes **True**.

Hence the sentence is **satisfiable**.

$$(\exists y \forall x (x = \text{Next}(y))) \rightarrow (\forall w \forall z (w = z))$$

Problem 2: Is it a tautology?

There is no “truth table method”.

You can't enumerate all possible interpretations!

$$(\exists y \forall x (x = \text{Next}(y))) \rightarrow (\forall w \forall z (w = z))$$

Problem 2: Is it a tautology?

Solution: Yes, it is a tautology!

Proof: Let I be any interpretation.

If $I[\exists y \forall x (x = \text{Next}(y))] = \mathbf{F}$,
then the sentence is **True**.

If $I[\exists y \forall x (x = \text{Next}(y))] = \mathbf{T}$,
then every object equals $\text{Next}(y)$.

In that case, $I[\forall w \forall z (w = z)] = \mathbf{T}$.

So no matter what, $I[\text{the sentence}] = \mathbf{T}$.

$$(\exists y \forall x (x = \text{Next}(y))) \rightarrow (\forall w \forall z (w = z))$$

Problem 2: Is it a tautology?

Is there **any** “mechanical method”??

More Inference Rules

\forall introduction

a var

..

..

P(a) true

$\forall x. P(x)$ true

\exists introduction

a var P(a) true

$\exists x. P(x)$ true

\exists elimination

a var P(a) true

..

..

$\exists x. P(x)$ true C true

C true

\forall elimination

a var $\forall x. P(x)$ true

P(a) true

Prove $\forall x. \exists y. y > x$ over natural number

x var

$x+1$ var

$\forall x. x+1 > x$ true

a var

a var

$\forall x. x+1 > x$ true

$a+1$ var

$a+1 > a$ true

\forall elimination

$\exists y. y > a$

\exists introduction

$\forall x. \exists y. y > x$

\forall introduction



Temporal logic

Propositional/FO logics have just one static state where formulae are evaluated

E.g.:

k stands for “it is snowing”

Is k true? No, but, only for today.

How to say:

It will snow someday in future.

It will snow everyday in future

Actually, it is possible to say the above in FOL, but, there is a much more elegant logic, which is also computationally easier to reason about

Temporal Logic Operators

Temporal operators:

Textual	Symbolic†	Explanation	Diagram
Unary operators:			
$X \phi$	$\bigcirc \phi$	neXt: ϕ has to hold at the next state.	
$G \phi$	$\square \phi$	Globally: ϕ has to hold on the entire subsequent path.	
$F \phi$	$\diamond \phi$	Finally: ϕ eventually has to hold (somewhere on the subsequent path).	
Binary operators:			
$\psi U \phi$	$\psi \mathcal{U} \phi$	Until: ψ has to hold at least until ϕ , which holds at the current or a future position.	
$\psi R \phi$	$\psi \mathcal{R} \phi$	Release: ϕ has to be true until and including the point where ψ first becomes true; if ψ never becomes true, ϕ must remain true forever.	

Textual	Symbolic†	Explanation	Diagram
Unary operators:			
$X \phi$	$\bigcirc \phi$	neXt: ϕ has to hold at the next state.	$\bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet$ ϕ is under the second state.
$G \phi$	$\square \phi$	Globally: ϕ has to hold on the entire subsequent path.	$\bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet$ ϕ is under each state.
$F \phi$	$\diamond \phi$	Finally: ϕ eventually has to hold (somewhere on the subsequent path).	$\bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet$ ϕ is under the third state.
Binary operators:			
$\psi U \phi$	$\psi \mathcal{U} \phi$	Until: ψ has to hold <i>at least</i> until ϕ , which holds at the current or a future position.	$\bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet$ ψ is under the first three states, ϕ is under the fourth.
$\psi R \phi$	$\psi \mathcal{R} \phi$	Release: ϕ has to be true until and including the point where ψ first becomes true; if ψ never becomes true, ϕ must remain true forever.	$\bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet$ ϕ is under the first three states, ϕ, ψ is under the fourth. $\bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet$ ϕ is under each state.


FG ϕ , GF ϕ ?

Temporal Logic with Past

F ("sometimes in the future"),
G ("always in the future"),
U ("until"), ...

P or F^{-1} for "once in the past",
H or G^{-1} for "always in the past",
S or U^{-1} for "since", ...


↓ $x \cdot \varphi$
At the current
time φ holds




Freeze Quantifier

$\downarrow x. \varphi$

At the current time φ holds



$\downarrow x. (\Diamond \downarrow y. ((x-y) \geq 14 \wedge \psi))$



Freeze Quantifier

Acknowledgement: Many Slides are from 15251